arXiv:1303.7219 KA–TP–07–2013 (v4)

arXiv:1303.7219v4 [gr-qc] 8 Apr 2013

Nonsingular spacetime defect

F.R. Klinkhamer* and C. Rahmede†

Institute for Theoretical Physics,
Karlsruhe Institute of Technology (KIT),
76128 Karlsruhe, Germany

Abstract

A nonsingular localized static classical solution is constructed for standard Einstein gravity coupled to an $SO(3) \times SO(3)$ chiral model of scalars [Skyrme model]. The construction proceeds in three steps. First, an *Ansatz* is presented for a solution with nontrivial topology of the spacetime manifold. Second, an exact vacuum solution of the reduced field equations is obtained. Third, matter fields are included and a particular numerical solution is found. This numerical solution has a negative effective mass, meaning that the gravitational force on a distant point mass is repulsive. Most likely, the origin of the negative effective mass traces back to the "surgery" needed to create the "defect" from Minkowski spacetime, but this process involves topology change and lies outside the realm of classical Einstein gravity.

PACS numbers: 04.20.Cv, 02.40.Pc

Keywords: general relativity, topology

---

*Electronic address: frans.klinkhamer@kit.edu

†Electronic address: christoph.rahmede@kit.edu

I. INTRODUCTION

It can be argued [1, 2] on general grounds that the small-length-scale structure of quantum spacetime is nontrivial. This structure has been called a quantum spacetime foam. Over larger length-scales (lower energy-scales), an effective classical spacetime manifold emerges and the crucial question is if that effective spacetime is perfectly smooth or not. Particle/wave propagation over Swiss-cheese-type manifolds has been studied and the problem can, in principle, be solved exactly [3, 4]. The simplest such Swiss-cheese-type manifold has identical static "defects" (alternatively called "holes" or "knots"), where each defect provides nontrivial topology. The particular defect considered has, however, a divergent (delta-function-type) Ricci curvature scalar and does not solve the vacuum Einstein equation.

The goal, now, is to construct a nonsingular defect solution by use of appropriate matter fields. For the defect topology at hand (holes with antipodal points identified; see below), it has been suggested [5] to use the gravitating $SO(3)$ Skyrme model [6, 7] with an appropriate interaction term [8] allowing for negative energy-density contributions. The present article uses a new *Ansatz* for the fields and gives special attention to the behavior of the reduced field equations at the defect core. We are, then, able to obtain a nonsingular defect solution in the gravitating $SO(3)$ Skyrme model.

Remarkably, the obtained solution displays the phenomenon of antigravity, i.e., a negative effective mass. As will become clear later on, precisely this antigravity property and the presence of negative energy densities allow a particular singularity theorem [9] to be circumvented.

II. MANIFOLD

The 4-dimensional spacetime manifold considered in this article is static for appropriate coordinates,

$$M_4 = \mathbb{R} \times M_3 \,. \tag{2.1}$$

The nontrivial topology appears in the 3-space M_3, which is, in fact, a noncompact, orientable, nonsimply-connected manifold without boundary. Up to a point, M_3 is homeomorphic to the 3-dimensional real-projective plane,

$$M_3 \simeq \mathbb{R}P^3 - \{\text{point}\} \,. \tag{2.2}$$

Further details can be found in Refs. [4, 5]. Here, only the necessary information is given.



Figure 1: Three-space M_3 obtained by surgery from $\mathbb{R}^3$: interior of ball with radius b removed and antipodal points on boundary of ball identified (as indicated by open and filled circles).

For the explicit construction of M_3, we perform some local surgery on the 3-dimensional Euclidean space $E_3 = \mathbb{R}^3$. We use the standard Cartesian and spherical coordinates on $\mathbb{R}^3$,

$$\vec{x} \equiv |\vec{x}|\, \widehat{x} = (x^1,\, x^2,\, x^3) = (r \sin\theta \cos\phi,\, r \sin\theta \sin\phi,\, r\cos\theta)\,, \tag{2.3}$$

with $x^m \in (-\infty,\, +\infty)$, $r \geq 0$, $\theta \in [0,\, \pi]$, and $\phi \in [0,\, 2\pi)$. Now, M_3 is obtained from $\mathbb{R}^3$ by the removal of the interior of the ball B_b with radius b and the identification of antipodal points on the boundary $S_b \equiv \partial B_b$. With point reflection denoted by $P(\vec{x}) = -\vec{x}$, the 3-space M_3 is given by

$$M_3 = \Big\{ \vec{x} \in \mathbb{R}^3 \;\Big|\; (|\vec{x}| \geq b > 0) \wedge (P(\vec{x}) \cong \vec{x} \;\text{ for }\; |\vec{x}| = b) \Big\}\,, \tag{2.4}$$

where $\cong$ stands for point-wise identification (Fig. 1).

The single set of coordinates (2.3) does not suffice for an appropriate description of M_3. The reason is simply that two different values of these coordinates may correspond to a single point. For example, $\vec{x} = (b,\, 0,\, 0)$ and $\vec{x} = (-b,\, 0,\, 0)$ correspond to really the same point of M_3. A relatively simple covering of M_3 uses three sets of coordinates (also called charts or patches), labeled by $n = 1, 2, 3$. Each coordinate patch surrounds one of the three Cartesian coordinate axes. The sets of coordinates are denoted

$$(X_n,\, Y_n,\, Z_n)\,, \quad \text{for} \quad n = 1,\, 2,\, 3\,, \tag{2.5}$$

which are, despite appearances, really non-Cartesian coordinates. Specifically, the set of coordinates surrounding the x^2-axis segment with $|x^2| \geq b$ is given by

$$X_2 \;=\; \begin{cases} \phi & \text{for} \quad 0 < \phi < \pi\,, \\ \phi - \pi & \text{for} \quad \pi < \phi < 2\pi\,, \end{cases} \tag{2.6a}$$

$$Y_2 \;=\; \begin{cases} r - b & \text{for} \quad 0 < \phi < \pi\,, \\ b - r & \text{for} \quad \pi < \phi < 2\pi\,, \end{cases} \tag{2.6b}$$

$$Z_2 \;=\; \begin{cases} \theta & \text{for} \quad 0 < \phi < \pi\,, \\ \pi - \theta & \text{for} \quad \pi < \phi < 2\pi\,, \end{cases} \tag{2.6c}$$

with ranges

$$X_2 \in (0,\, \pi)\,, \tag{2.7a}$$

$$Y_2 \in (-\infty,\, \infty)\,, \tag{2.7b}$$

$$Z_2 \in (0,\, \pi)\,. \tag{2.7c}$$

The other two sets, $(X_1,\, Y_1,\, Z_1)$ and $(X_3,\, Y_3,\, Z_3)$, are defined similarly.

In the following, we consider spherically symmetric fields and it suffices to consider one set of coordinates, which we take to be $n = 2$ set (2.6). The notation is simplified as follows:

$$(X,\, Y,\, Z,\, T) \equiv (X_2,\, Y_2,\, Z_2,\, t)\,, \tag{2.8}$$

where the time coordinate has been added in order to describe the spacetime manifold M_4.

III. FIELDS AND ACTION

The spacetime manifold (2.1) of the previous section is now equipped with a metric $g_{\mu\nu}$, whose dynamics is taken to be governed by the standard Einstein–Hilbert action [10]. In addition, there is assumed to be a scalar field $\Omega \in SO(3)$, with self-interactions governed by a Skyrme term in the action [6] and by another term [8] whose coupling constant γ is taken to be positive, allowing for negative energy-density contributions.

Specifically, the combined action of the pure-gravity sector, labeled 'grav,' and the matter sector, labeled 'mat,' is given by $(c = \hbar = 1)$

$$S = \int_{M_4} d^4X\, \sqrt{-g}\, \Big(\mathcal{L}_{\text{grav, EH}} + \mathcal{L}_{\text{mat, kin}} + \mathcal{L}_{\text{mat, Skyrme}} + \mathcal{L}_{\text{mat, metastab}} \Big)\,, \tag{3.1a}$$

where

$$\mathcal{L}_{\text{grav, EH}} = \frac{1}{16\pi G_N}\, R\,, \tag{3.1b}$$

$$\mathcal{L}_{\text{mat, kin}} = \frac{f^2}{4}\, \text{tr}\big(\omega_\mu\, \omega^\mu\big)\,, \tag{3.1c}$$

$$\mathcal{L}_{\text{mat, Skyrme}} = \frac{1}{16e^2}\, \text{tr}\Big([\omega_\mu,\, \omega_\nu]\, [\omega^\mu,\, \omega^\nu] \Big)\,, \tag{3.1d}$$

$$\mathcal{L}_{\text{mat, metastab}} = \gamma\, \frac{1}{48e^2}\, \Big(\text{tr}(\omega_\mu\, \omega^\mu)\Big)^2\,, \tag{3.1e}$$

in terms of an $so(3)$-Lie-algebra valued one-form $\omega \equiv \omega_\mu\, dX^\mu$ given by

$$\omega_\mu \equiv \Omega^{-1} \partial_\mu \Omega\,. \tag{3.1f}$$

The $SO(3) \times SO(3)$ global symmetry of the matter sector is realized on the scalar field as follows:

$$\Omega \to S_L \cdot \Omega \cdot S_R^{-1}\,, \tag{3.2}$$

where S_L, $S_R \in SO(3)$ and the central dot denotes matrix multiplication. As mentioned at the end of Sec. II, the measure d^4X in the integral (3.1a) corresponds to only one of the three coordinate charts needed to cover M_4.

IV. ANSATZ AND FIELD EQUATIONS

A. Ansatz

A spherically symmetric *Ansatz* for the metric is given by the following line element:

$$ds^2 = -e^{2\widetilde{\nu}(W)}\,dT^2 + e^{2\widetilde{\lambda}(W)}\,dY^2 + W\left(dZ^2 + \sin^2 Z\,dX^2\right)\,, \tag{4.1a}$$

$$W \equiv b^2 + Y^2\,, \tag{4.1b}$$

where, as explained at the end of Sec. II, we only show the coordinates of one chart with $Y \in (-\infty, \infty)$. The scalar field is given by the hedgehog-type *Ansatz* [5, 6],

$$\Omega = \cos\left[\widetilde{F}\left(r^2\right)\right]\,\mathbb{1} - \sin\left[\widetilde{F}\left(r^2\right)\right]\,\widehat{x}\cdot\vec{S} + \left(1 - \cos\left[\widetilde{F}\left(r^2\right)\right]\right)\,\widehat{x}\otimes\widehat{x}\,, \tag{4.2a}$$

$$\widetilde{F}(b^2) = \pi\,, \tag{4.2b}$$

$$S_1 \equiv \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \quad S_2 \equiv \begin{pmatrix} 0 & 0 & -1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad S_3 \equiv \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \tag{4.2c}$$

where, in components, $(\widehat{x}\otimes\widehat{x})^{ab} = \widehat{x}^a\,\widehat{x}^b$. Remark that, because of the boundary condition (4.2b) at $|\vec{x}| = b$, it is possible to use the single coordinate chart (2.3) with the further identification $r^2 = b^2 + Y^2 = W$ for the coordinates used in the metric (4.1). In other words, the topology of M_3 (see Fig. 1) is trivially consistent with the hedgehog field having boundary value $\widetilde{F}(b^2) = \pi$.

The arguments of the tilde-functions in the above *Ansätze* have dimension length-square. Related functions with lengths as arguments can be defined as follows:

$$\nu\left(\sqrt{b^2 + Y^2}\right) \equiv \widetilde{\nu}\left(b^2 + Y^2\right)\,, \tag{4.3}$$

and similarly for λ and F. The functions without tilde resemble those of the previous literature [6–8], but we prefer to work with the tilde-functions.

The relevant nonvanishing components of the Riemann tensor for the *Ansatz* (4.1) are:

$$R^{T}{}_{YTY} = -2\left[\widetilde{\nu}' + 2Y^2\left(-\widetilde{\nu}'\,\widetilde{\kappa}'/\widetilde{\kappa} + \widetilde{\nu}'' + \widetilde{\nu}'^2\right)\right], \tag{4.4a}$$

$$R^{T}{}_{ZTZ} = -2Y^2\,\widetilde{\nu}'/\widetilde{\kappa}^2, \tag{4.4b}$$

$$R^{T}{}_{XTX} = \sin^2 Z\,R^{T}{}_{ZTZ}, \tag{4.4c}$$

$$R^{Y}{}_{ZYZ} = \frac{2Y^2\,(b^2+Y^2)\,\widetilde{\kappa}'/\widetilde{\kappa} - b^2}{(b^2+Y^2)\,\widetilde{\kappa}^2}, \tag{4.4d}$$

$$R^{Y}{}_{XYX} = \sin^2 Z\,R^{Y}{}_{ZYZ}, \tag{4.4e}$$

$$R^{Z}{}_{XZX} = \sin^2 Z\left(1 - \frac{Y^2}{(b^2+Y^2)\,\widetilde{\kappa}^2}\right), \tag{4.4f}$$

where the prime stands for differentiation with respect to W. The components not shown in (4.4) either vanish or can be computed from the ones above by the usual symmetry properties.

B. Reduced field equations

The derivation of the variational equations is straightforward.[1] Henceforth, we will use the following dimensionless model parameters and dimensionless variables:

$$\widetilde{\eta} \equiv 8\pi\eta \equiv 8\pi\,G_N\,f^2, \tag{4.5a}$$

$$w \equiv (e\,f)^2\,W = (y_0)^2 + y^2, \tag{4.5b}$$

$$y \equiv e\,f\,Y, \tag{4.5c}$$

$$y_0 \equiv e\,f\,b. \tag{4.5d}$$

Furthermore, the following functions are introduced:

$$\widetilde{\kappa}(w) \equiv \exp\left[\widetilde{\lambda}(w)\right], \tag{4.6a}$$

$$\widetilde{\mu}(w) \equiv \exp\left[\widetilde{\nu}(w)\right]. \tag{4.6b}$$

The reduced Einstein equations and matter field equation will be given in Appendix A.

---

[1] The variational equations for $\gamma = 0$ can also be obtained by making appropriate rescalings ($F \to F/2$ and $e^2 \to e^2/2$) and substitutions ($\nu \to \widetilde{\nu}$, $\lambda \to \widetilde{\lambda}$, and $F \to \widetilde{F}$) in the equations of Ref. [7].

From these equations, one obtains three ordinary differential equations (ODEs):

$$\widetilde{\nu}'(w) \;=\; \frac{\widetilde{\kappa}(w)^2}{4\,(w-y_0^2)} - \frac{1}{4w} + \widetilde{\eta}\,Q_1\big[\widetilde{F}(w),\, \widetilde{\kappa}(w),\, \widetilde{\nu}(w)\big]\,, \tag{4.7a}$$

$$\widetilde{\kappa}'(w) \;=\; \widetilde{\kappa}(w)\,\frac{w+y_0^2-\widetilde{\kappa}(w)^2}{4w\,(w-y_0^2)} + \widetilde{\eta}\,Q_2\big[\widetilde{F}(w),\, \widetilde{\kappa}(w),\, \widetilde{\nu}(w)\big]\,, \tag{4.7b}$$

$$\widetilde{F}''(w) \;=\; Q_3[\widetilde{F}(w),\, \widetilde{\kappa}(w),\, \widetilde{\nu}(w)]\,, \tag{4.7c}$$

where the prime now stands for differentiation with respect to w and Q_n are certain functionals given in Appendix A. These ODEs are to be solved with the following boundary conditions:

$$\widetilde{F}(0) \;=\; \widetilde{F}(\infty) = \pi\,, \tag{4.8a}$$

$$\widetilde{\kappa}(0) \;=\; 0\,, \quad \widetilde{\nu}(\infty) = 0\,. \tag{4.8b}$$

The field configuration (4.2) with boundary conditions (4.8) has winding number zero [recall (2.2) and the further homeomorphism $SO(3) \simeq \mathbb{R}P^3$]. It is, therefore, not a genuine Skyrmion [which has $\widetilde{F}(\infty) = 0$ and winding number one]. Our matter solution may very well turn out to be unstable, but that is not important for one of the main goals of this article, i.e., finding a possible origin of antigravity effects (see Sec. V).

C. Power-series solution

An analytic solution near the defect core can be obtained as a power series. The first terms are as follows:

$$F(w) \;=\; \pi - f\,(w-y_0^2) + \cdots\,, \tag{4.9a}$$

$$\widetilde{\kappa}(w) \;=\; k\,\sqrt{w-y_0^2} + \cdots\,, \tag{4.9b}$$

$$\widetilde{\nu}(w) \;=\; \widetilde{\nu}_0 - n\,(w-y_0^2) + \cdots\,, \tag{4.9c}$$

with coefficients f, k, and n, and the arbitrary constant $\widetilde{\nu}_0$. Given the value for k, the reduced field equations fix the values (denoted by an overbar) of the other two constants:

$$\overline{f} \;=\; \sqrt{\frac{k^2\;(12-8\,\gamma+3\,{y_0}^2)}{4\,\gamma\,{y_0}^2}}\,, \tag{4.10a}$$

$$\overline{n} \;=\; k^2\left[-\frac{1}{4} + 3\,\widetilde{\eta} - \frac{3\,\widetilde{\eta}}{\gamma} + \frac{9\,\widetilde{\eta}}{{y_0}^2} - \frac{6\,\widetilde{\eta}}{\gamma\,{y_0}^2} - \frac{4\,\widetilde{\eta}\,\gamma}{{y_0}^2} - \frac{3\,\widetilde{\eta}\,{y_0}^2}{8\,\gamma}\right]\,, \tag{4.10b}$$

where γ is assumed to be positive and not too large, so that $12-8\,\gamma+3\,{y_0}^2 \geq 0$.

With the leading terms of the above power-series solutions, it is easy to verify that the components of the Riemann tensor (4.4) are finite at the defect core $y = 0$.

V. EXACT VACUUM SOLUTION

For vacuum matter fields,

$$\widetilde{F}(w) \;=\; 0\,, \tag{5.1a}$$

the ODEs (4.7a) and (4.7b) have $Q_1 = Q_2 = 0$ and can be solved exactly for boundary conditions (4.8b),

$$\widetilde{\kappa}(w) \;\equiv\; \exp\left[\widetilde{\lambda}(w)\right] = \sqrt{\frac{1-(y_0)^2/w}{1-\ell/\sqrt{w}}}\;, \tag{5.1b}$$

$$\widetilde{\mu}(w) \;\equiv\; \exp\left[\widetilde{\nu}(w)\right] = \sqrt{1-\ell/\sqrt{w}}\;, \tag{5.1c}$$

with a dimensionless constant ℓ and the definition $w \equiv y^2 + (y_0)^2$ in terms of the 'radial' coordinate $y \in (-\infty,\, +\infty)$ and the defect parameter $y_0 > 0$ (the corresponding dimensional parameter b is also nonzero and positive). The same solution is obtained for the case $\widetilde{F}(w) = \pi$ and $\widetilde{\eta} = 0$.

At this point, we can already make two general remarks. First, the vanishing of the metric component $\widetilde{\kappa}^2$ at $Y = 0$ implies that the defect of Fig. 1 at a fixed time T has zero physical extent along the radial direction.

Second, the real $\widetilde{\kappa}$ and $\widetilde{\mu}$ functions from (5.1), for large enough values of $|\ell|$, cover the whole of the manifold M_4 only if ℓ is $\underline{\text{negative}}$:

$$\ell^2 > (y_0)^2 \quad\Rightarrow\quad \ell < -y_0 < 0\,. \tag{5.2}$$

In turn, this implies having $\underline{\text{anti}}$-gravity, because asymptotically ($w \sim y^2$) the solution behaves as follows:

$$\widetilde{\kappa}^2 \;\sim\; 1/\big(1-\ell/|y|\big)\,, \tag{5.3a}$$

$$\widetilde{\mu}^2 \;\sim\; \big(1-\ell/|y|\big)\,, \tag{5.3b}$$

which is to be compared to the standard Schwarzschild metric [10] with positive mass M, having the respective components $1/(1-2G_N M/r)$ and $(1-2G_N M/r)$. For the vacuum solution (5.1) with $\ell < 0$, a point mass far away from the defect core will not be attracted towards it but repulsed.

For $\ell < y_0$, all Riemann-curvature-tensor components (4.4) are finite over the whole manifold, also at the defect core, $y = 0$ or $w = (y_0)^2$. Specifically, we find for the relevant nonvanishing Riemann tensor components:

$$R^{T}{}_{YTY} \;=\; (e^2 f^2)\,\frac{\ell\,(w-y_0^2)}{w^2\,(\sqrt{w}-\ell)}\,, \tag{5.4a}$$

$$R^{T}{}_{ZTZ} \;=\; \frac{-\ell}{2\sqrt{w}}\,, \tag{5.4b}$$

$$R^{Y}{}_{ZYZ} \;=\; \frac{-\ell}{2\sqrt{w}}\,, \tag{5.4c}$$

$$R^{Z}{}_{XZX} \;=\; \sin^2 Z\,\frac{\ell}{\sqrt{w}}\,. \tag{5.4d}$$

More importantly and for all values of ℓ, the scalar obtained by contraction of the Riemann tensor with itself,

$$S_2 \;\equiv\; R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} = 12\,\frac{\ell^2}{w^3}\,, \tag{5.4e}$$

remains finite over the whole of the manifold M_4, because $w \geq (y_0)^2 > 0$. This behavior contrasts with that of the Schwarzschild metric over $\mathbb{R}^4$, for which S_2 diverges at the point $r = 0$.[2]

A brief discussion of the geodesics from the metric (4.1) with functions (5.1b) and (5.1c) is given in Appendix B. A more extensive discussion of geodesics is scheduled for a follow-up paper, while another follow-up paper intends to discuss the global structure of the new vacuum solution over M_4.[3]

Note, finally, that the regular flat-spacetime metric is given by (5.1b) and (5.1c) with $\ell = 0$. The actual value of the free parameter ℓ in the metric from (5.1) will have to be determined by adding matter fields (the same applies for the determination of M in the standard Schwarzschild solution). An interesting question is if it is possible to get negative ℓ, corresponding to antigravity.

VI. NUMERICAL SOLUTION WITH MATTER FIELDS

We use the following values for the model parameter γ and the defect parameter y_0:

$$\overline{\gamma} \;=\; 3/2 + \Delta\,, \quad \Delta \geq 0\,, \tag{6.1a}$$

$$\overline{y_0} \;=\; \sqrt{\big[\overline{\gamma}\,(8+\delta^2) - 12\big]/3}\,. \tag{6.1b}$$

---

[2] With $0 < b \ll \ell/(ef) = 2G_N M$, the manifold (2.1) and metric (4.1), (5.1b), (5.1c) provide a "regularization" of the standard Schwarzschild solution.

[3] Electromagnetic wave propagation over this spacetime can be calculated with the methods of Refs. [3, 4].

These parameters for $\delta \to 0$ are consistent with having zero coefficient f and positive finite coefficient k in the series solution (4.9), according to (4.10). The numerical solution of the ODEs will be obtained for $w \in [w_{\text{min}},\, w_{\text{max}}]$, with $w_{\text{min}} = (\overline{y_0})^2 + \epsilon^2$ and large enough w_{max}. The boundary conditions at $w_{\text{bcs}} = w_{\text{min}}$ follow from the analytic expressions (4.9). Anticipating the discussion of Sec. VIII, the coupling constant (4.5a) is taken infinitesimal, $\eta = \epsilon^2$.

For parameters (6.1) with $\delta = 0$ and $\Delta = 1/100$ and for a particular value of the coefficient k, the numerical solution is given in Figs. 2 and 3. The dimensionless energy density is seen to be negative at the defect core $\big[$with $t_{00,\,\text{def}} = 4\, e^{2\widetilde{\nu}_0}\,\big(1 + y_0^2 - 4\gamma/3\big)\big/(y_0)^4\,\big]$ and to have a positive tail with diverging total energy [see Sec. VIII for further discussion].

From the asymptotic behavior of the functions shown in Fig. 2, the parameter ℓ as defined in (5.3) is found to take the value

$$\ell \approx -28\,, \tag{6.2}$$

also for smaller values of ϵ. The numerical result (6.2) suggests that it is possible to realize the vacuum solution (5.1) with a negative mass parameter ℓ. Still, the numerical analysis reported in this article is only exploratory and needs to be confirmed.

VII. DEFECT MASS

According to (5.3), the parameter ℓ of the vacuum solution (5.1) is proportional to twice the Schwarzschild mass (see Sec. VIII for further discussion of the energy scales of the model). Another useful definition of the mass is the so-called Komar mass [10, 11]. For the *Ansatz* (4.1), the dimensionless Komar mass at infinity becomes

$$\widehat{m}_{\text{Komar}} = \lim_{\widehat{w}\to\infty} m_{\text{Komar}}(\widehat{w}) = \lim_{\widehat{w}\to\infty} \frac{\widetilde{\mu}(\widehat{w})}{\widetilde{\kappa}(\widehat{w})}\, 2\widehat{w}\, \sqrt{\widehat{w} - y_0^2}\; \widetilde{\nu}'(\widehat{w})\,. \tag{7.1}$$

This mass comes from an integral over a distant sphere (having radius square $w = \widehat{w}$), but, for the vacuum solution, it can be transformed [5] into an integral over the defect sphere (having $w = y_0^2$) with the result:

$$\widehat{m}_{\text{Komar}} = m_{\text{Komar}}(y_0^2) = \ell/2\,. \tag{7.2}$$

Even with matter fields included, as was the case for the numerical solution of Sec. VI, the negative contribution to the Komar-mass from the "defect" at $r = b$ can still dominate the standard (positive) contributions from gravity and matter in the bulk. Precisely this net antigravity property and the presence of negative energy density at the defect core invalidate



Figure 2: Numerical solution of the ODEs (A4) over the w interval $\{w_{\rm min},\, w_{\rm max}\} = \{2/75 + \epsilon^2, 10^6\}$ with $\epsilon = 10^{-4}$ and model parameters $\{\gamma,\, \eta,\, y_0\} = \{151/100,\, 10^{-8},\, \sqrt{2/75}\}$. The boundary conditions at $w_{\rm bcs} = w_{\rm min} = y_0^2 + \epsilon^2$ follow from the analytic expressions (4.9) with coefficients $\{k,\, n,\, f\} = \{\sqrt{8/75},\, -\big(1/4+37\,\pi/12500000\big)\, 8/75,\, 0\}$ and constant $\widetilde{\nu}_0$ adjusted to give $\widetilde{\nu}(w_{\rm max}) = -\widetilde{\lambda}(w_{\rm max})$.



Figure 3: Dimensionless energy density $t_{00} \equiv T_{00}/(e^2\, f^4)$ for the functions from Fig. 2.

the Singularity Theorem 2.1 of Gannon [9], which applies to spacetimes with asymptotically-flat nonsimply-connected spacelike slices. This suggests that having a nonsingular defect solution may result in a negative mass, also for matter sectors which are different from the particular Skyrme model considered here.

The defect mass discussed up till now is the gravitational mass. It remains to be seen what the inertial properties of the defect are.

VIII. DISCUSSION

The numerical solution from Sec. VI has asymptotically a nonzero energy density, (Fig. 3)

$$T_{00} \propto \frac{1}{w}\, e^2 f^4 \sim \frac{f^2}{r^2}\,, \tag{8.1}$$

where we have used $w \sim y^2 \sim (efr)^2$ in terms of the standard radial coordinate r and the parameters e and f from the action (3.1). The asymptotic value of the Komar mass of the numerical solution is

$$\widehat{M}_{\text{Komar}} \sim \frac{1}{e\,f\,G_N}\,\ell/2\,, \tag{8.2}$$

where the coefficient ℓ is obtained from the behavior $2\widetilde{\lambda} \sim -2\widetilde{\nu} \sim -\ell/|y|$ in (5.3).

Of interest to physics are, therefore, the following mathematical limits:

$$f \;\to\; 0\,, \tag{8.3a}$$

$$e \;\to\; \infty\,, \tag{8.3b}$$

$$e\,f \;\to\; \text{const} > 0\,, \tag{8.3c}$$

while G_N is kept fixed (recall $c = 1$). The limit (8.3a) makes for vanishing energy density (8.1), whereas the limit (8.3c) keeps the Komar mass (8.2) finite.

The numerical solution of Fig. 2 has a Schwarzschild-like behavior (5.3) with $\ell < 0$. The corresponding nonsingular defect in the limits (8.3) then produces a finite negative mass, i.e., antigravity, while providing for a negligible local energy density (the total scalar-field energy up to a fixed physical distance also vanishes). However, as mentioned at the end of Sec. VI, the numerical analysis of the present article still needs to be verified and extended.

Note that, strictly speaking, the $SO(3)$ scalar field is absent in the Standard Model. But the nonlinear sigma model resurfaces if the gauge fields are eliminated and the Higgs-field modulus is frozen. Hence, the toy model of matter fields considered in the present article is not completely removed from realistic physics. As such, there may even be a connection with ideas linking the quantum structure of spacetime (having an energy scale $E_{\text{Planck}} \equiv \sqrt{\hbar\, c^5/G_N}$ and a length scale $\hbar c/E_{\text{Planck}}$) to the Higgs-boson and top-quark masses [12]. We refrain from making even wilder speculations about the cosmological constant. In fact, the main focus of this article remains purely theoretical, namely, finding a nonsingular spacetime-defect solution and a possible mechanism for the origin of antigravity effects.

ACKNOWLEDGMENTS

The authors acknowledge support by the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG) under grant number KL 1103/2-1.

Appendix A: ODEs

In this appendix, the relevant ordinary differential equations (ODEs) are given in detail, whereas they were only given in abbreviated form in (4.7). It turns out to be helpful to define the following dimensionless auxiliaries:

$$u_1 \equiv \left(1-\frac{4}{3}\,\gamma\right)\sin^2\left(\frac{\widetilde{F}}{2}\right), \tag{A1a}$$

$$u_2 \equiv \left(1-\frac{2}{3}\,\gamma\right)\sin^2\left(\frac{\widetilde{F}}{2}\right)+\frac{w}{4}, \tag{A1b}$$

$$u_3 \equiv \widetilde{\eta}\left[\left(1-\frac{4}{3}\,\gamma\right)\cos(2\widetilde{F})-4\left(1-\frac{4}{3}\,\gamma+w\right)\cos\widetilde{F}+3\left(1-\frac{4}{3}\,\gamma+\frac{4}{3}\,w\right)\right]-2w\,. \tag{A1c}$$

The reduced Einstein equations can now be written in the form ${G^{Y}}_{Y}=8\pi G_N {T^{Y}}_{Y}$, ${G^{Y}}_{Y}-{G^{T}}_{T}=8\pi G_N({T^{Y}}_{Y}-{T^{T}}_{T})$, and ${G^{Z}}_{Z}=8\pi G_N {T^{Z}}_{Z}$ [it is also found that ${G^{X}}_{X}={G^{Z}}_{Z}$, ${T^{X}}_{X}={T^{Z}}_{Z}$]. In terms of dimensionless quantities, these reduced Einstein equations are:

$$\begin{aligned}&1-\frac{(w-y_0^2)\,e^{-2\widetilde{\lambda}}\,(4w\,\widetilde{\nu}'+1)}{w}=\\&4\widetilde{\eta}\left[\left(1+\frac{u_1}{w}\right)\sin^2\left(\frac{\widetilde{F}}{2}\right)+\left(w-y_0^2\right)e^{-2\widetilde{\lambda}}\,\widetilde{F}'^2\left(-2u_2+\gamma w\left(w-y_0^2\right)e^{-2\widetilde{\lambda}}\,\widetilde{F}'^2\right)\right],\end{aligned} \tag{A2a}$$

$$\begin{aligned}&2\left(w-y_0^2\right)\left(\widetilde{\lambda}'+\widetilde{\nu}'\right)-\frac{y_0^2}{w}=\\&8\widetilde{\eta}\left(w-y_0^2\right)\widetilde{F}'^2\left(u_2-\frac{1}{3}\gamma w\left(w-y_0^2\right)e^{-2\widetilde{\lambda}}\,\widetilde{F}'^2\right),\end{aligned} \tag{A2b}$$

$$\begin{aligned}&-e^{-2\widetilde{\lambda}}\left(2w\left(w-y_0^2\right)\left(\widetilde{\lambda}'\,\widetilde{\nu}'-\widetilde{\nu}'^2-\widetilde{\nu}''\right)+\left(w-y_0^2\right)\widetilde{\lambda}'-\left(2w-y_0^2\right)\widetilde{\nu}'-\frac{y_0^2}{2w}\right)=\\&\frac{2}{w}\,\widetilde{\eta}\,u_1\sin^2\left(\frac{\widetilde{F}}{2}\right)-w\,\widetilde{\eta}\left(w-y_0^2\right)e^{-2\widetilde{\lambda}}\,\widetilde{F}'^2\left(1-\frac{2}{3}\,\gamma\left(w-y_0^2\right)e^{-2\widetilde{\lambda}}\,\widetilde{F}'^2\right).\end{aligned} \tag{A2c}$$

The reduced matter field equation is

$$\begin{aligned}
0 \;=\; & \left(1+\frac{2u_1}{w}\right) e^{2\widetilde{\lambda}} \sin\left(\widetilde{F}\right) \\
& -16\sqrt{w-z_0^2}\,\widetilde{F}' \left[u_2\left(\left(w-z_0^2\right)\left(\widetilde{\nu}'-\widetilde{\lambda}'\right)+1\right)+\frac{1}{4}\left(w-z_0^2\right)\right] \\
& -\frac{16}{3}\gamma\left(w-z_0^2\right)^{3/2} e^{-2\widetilde{\lambda}}\,\widetilde{F}'^3 \left[w\left(w-z_0^2\right)\left(3\widetilde{\lambda}'-\widetilde{\nu}'\right)-3w+z_0^2\right] \\
& +2\left(1-\frac{2\gamma}{3}\right)\left(w-z_0^2\right)\left(1-4\sqrt{w-z_0^2}\right)\widetilde{F}'^2 \sin\widetilde{F} \\
& -16\left(w-z_0^2\right)^{3/2}\widetilde{F}''\left(u_2-\gamma w\left(w-z_0^2\right)e^{-2\widetilde{\lambda}}\,\widetilde{F}'^2\right). \qquad \text{(A3)}
\end{aligned}$$

Using $\widetilde{\kappa}(w)$ as defined in (4.6a) and solving for $\widetilde{\nu}'$, $\widetilde{\kappa}'$, $\widetilde{F}''$, one obtains the final ODEs:

$$\begin{aligned}
\widetilde{\nu}' \;=\; & \frac{\widetilde{\kappa}^2}{4\left(w-y_0^2\right)}-\frac{1}{4w}-\frac{\widetilde{\eta}\,\widetilde{\kappa}^2}{w\left(w-y_0^2\right)}\sin^2\left(\frac{\widetilde{F}}{2}\right)\left(u_1+w\right)+2\widetilde{\eta}\,u_2\,\widetilde{F}'^2 \\
& -\gamma\widetilde{\eta}w\left(w-y_0^2\right)\widetilde{\kappa}^{-2}\,\widetilde{F}'^4\,, \qquad \text{(A4a)}
\end{aligned}$$

$$\widetilde{\kappa}' \;=\; \widetilde{\kappa}\,\frac{w+y_0^2-\widetilde{\kappa}^2}{4w\left(w-y_0^2\right)}+2\widetilde{\eta}\,u_2\,\widetilde{F}'^2-\frac{\gamma}{3}\widetilde{\eta}w\left(w-y_0^2\right)\widetilde{\kappa}^{-2}\,\widetilde{F}'^4\,, \qquad \text{(A4b)}$$

$$\begin{aligned}
\widetilde{F}'' \;=\; & \frac{1}{24\left(u_2-\gamma w\left(w-y_0^2\right)\widetilde{\kappa}^{-2}\,\widetilde{F}'^2\right)} \\
& \left[\frac{3\widetilde{\kappa}^2}{w\left(w-y_0^2\right)^{3/2}}\left(u_1+\frac{w}{2}\right)\sin\widetilde{F}-\frac{3}{w-y_0^2}\left(-\frac{2}{w}\,\widetilde{\kappa}^2\,u_2u_3+4u_2+2w-2y_0^2\right)\widetilde{F}'\right. \\
& +\frac{3}{\sqrt{w-y_0^2}}\left(1-4\sqrt{w-y_0^2}\right)\left(1-\frac{2}{3}\gamma\right)\sin\widetilde{F}\;\widetilde{F}'^2 \\
& \left.-4\gamma\left(u_3+\left(-4w+3y_0^2\right)\widetilde{\kappa}^{-2}\right)\widetilde{F}'^3-16\gamma\,\widetilde{\eta}\,w\left(w-y_0^2\right)\,\widetilde{\kappa}^{-2}\,u_2\,\widetilde{F}'^5\right]. \qquad \text{(A4c)}
\end{aligned}$$

Appendix B: Vacuum-solution geodesics

For the vacuum solution of Sec. V, the nontrivial Christoffel symbols are

$$\frac{1}{ef}\Gamma^{T}{}_{TY} \;=\; \frac{\sqrt{w-y_0^2}\;\ell/2}{w^{3/2}-\ell\,w}\,, \qquad \text{(B1a)}$$

$$\frac{1}{ef}\Gamma^{Y}{}_{TT} \;=\; \frac{\left(\sqrt{w}-\ell\right)\,\ell/2}{w\sqrt{w-y_0^2}}\,, \qquad \text{(B1b)}$$

$$\frac{1}{ef}\Gamma^{Y}{}_{YY} \;=\; \frac{\sqrt{w}\,y_0^2-\left(w+y_0^2\right)\,\ell/2}{w\left(\sqrt{w}-\ell\right)\sqrt{w-y_0^2}}\,. \qquad \text{(B1c)}$$

The last two Christoffel symbols are seen to diverge at the defect core ($w = y_0^2$), but the particle motion can still be regular, as we will now show by a simple example [using dimensionless Christoffel symbols $\gamma^a_{bc} \equiv (ef)^{-1}\,\Gamma^A_{BC}$].

Consider, in fact, the geodesic equation for a particle moving solely in the y-direction and denote the nonvanishing dimensionless velocity components $u^t = dt/d\tau$ and $u^y = dy/d\tau$, where τ is the dimensionless proper time and $c = 1$. The particle motion is, then, given by the following two equations:

$$0 \;=\; \frac{du^t}{d\tau} + 2\,\gamma^t_{\;ty}\,u^t u^y\,, \tag{B2a}$$

$$0 \;=\; \frac{du^y}{d\tau} + \gamma^y_{\;tt}\,u^t u^t + \gamma^y_{\;yy}\,u^y u^y\,. \tag{B2b}$$

The first equation can be solved for u^t:

$$u^t = \left(1 - \frac{\ell}{\sqrt{w}}\right)^{-1}\,. \tag{B3}$$

Inserting this u^t into the second equation gives

$$\frac{d^2 y}{d\tau^2} + \frac{-y^2\,\ell + 2y_0^2\left(-\ell + \sqrt{y_0^2+y^2}\right)}{2y\,(y^2+y_0^2)\left(-\ell + \sqrt{y_0^2+y^2}\right)}\left(\frac{dy}{d\tau}\right)^2 = \frac{\ell}{2y\left(\ell - \sqrt{y_0^2+y^2}\right)}\,. \tag{B4}$$

Finally, replace in (B4)

$$\frac{dy}{d\tau} = \frac{1}{2y}\frac{dy^2}{d\tau}\,,\quad \frac{d^2y}{d\tau^2} = \frac{1}{2y}\frac{d^2y^2}{d\tau^2} - \frac{1}{4y^3}\left(\frac{dy^2}{d\tau}\right)^2 \tag{B5}$$

to obtain

$$\frac{d^2y^2}{d\tau^2} - \frac{1}{4(y_0^2+y^2)}\left(1 + \frac{\sqrt{y_0^2+y^2}}{-\ell + \sqrt{y_0^2+y^2}}\right)\left(\frac{dy^2}{d\tau}\right)^2 = \frac{\ell}{\ell - \sqrt{y_0^2+y^2}}\,, \tag{B6}$$

which remains finite along the trajectory as long as $\ell < y_0$.

---

[1] J.A. Wheeler, "On the nature of quantum geometrodynamics," Ann. Phys. (N.Y.) **2**, 604 (1957).

[2] J.A. Wheeler, "Superspace and the nature of quantum geometrodynamics," in: *Battelle Rencontres 1967*, edited by C.M. DeWitt and J.A. Wheeler (Benjamin, New York, USA, 1968), Chap. 9.

[3] H.A. Bethe, "Theory of diffraction by small holes," Phys. Rev. **66**, 163 (1944).

[4] S. Bernadotte and F.R. Klinkhamer, “Bounds on length scales of classical spacetime foam models,” Phys. Rev. D **75**, 024028 (2007); arXiv:hep-ph/0610216.

[5] M. Schwarz, “Nontrivial spacetime topology, modified dispersion relations, and an $SO(3)$-Skyrme model, PhD Thesis, KIT, July 2010.

[6] T.H.R. Skyrme, “A nonlinear field theory,” Proc. Roy. Soc. Lond. A **260**, 127 (1961).

[7] N.K. Glendenning, T. Kodama, and F.R. Klinkhamer, “Skyrme topological soliton coupled to gravity,” Phys. Rev. D **38**, 3226 (1988).

[8] D.E.L. Pottinger and E. Rathske, “Metastability of solitons in a generalized Skyrme model,” Phys. Rev. D **33**, 2448 (1986).

[9] D. Gannon, “Singularities in nonsimply connected space-times,” J. Math. Phys. **16**, 2364 (1975).

[10] R.M. Wald, *General Relativity* (Chicago University Press, 1984).

[11] A. Komar, “Covariant conservation laws in general relativity,” Phys. Rev. **113**, 934 (1959).

[12] F.R. Klinkhamer, “Standard Model Higgs field and energy scale of gravity,” to appear in JETPL, arXiv:1302.1496.